MEDICATION INFORMATION

*Use weight to determine dose especially if child was premature or is small for his/her age.

MEDICATIONS

MEDICATIONS					
Age	1-3 months	4-6 months	7-12 months	13-24 months	2-3 years
Weight	7-12 lbs	13-18 lbs	19-24 lbs	25-30 lbs	31-38 lbs
	<u>4 ml</u>	¹/2 tsp	3∕4 tsp	1 tsp	1 ½ tsp
	<u>¹⁄₄ tsp</u>	¹/2 tsp	3/4 tsp	1 tsp	1 ½ tsp
Acetaminophen (Tylenol, Tempra, etc.)	4 ml	8 ml	1.2 ml	1.6 ml	2.0 ml
Every 4 hours as needed For fever or pain. Drops In ml, elixir in tsp. and	<u>½ tsp</u>	½ tsp	3/4 tsp	1 tsp	1 ½ tsp
Chewable in mgs. Instead					200 mgs
Ibuprofen (Advil, Motrin)			27.4		2 ½ mgs
Every 6 hours, may be give	en		<u>34 tsp</u>	1 tsp	1 ½ tsp
Pedicare infant drops Triaminic infant or	(check w/dr.)	8 ml	1.2 ml	1.6 ml	2.0 ml
Pediatric Sudafed Drops Every 6 hours as needed For nasal congestion					
Benadryl Elixir Every 4-6 hours for Allergic symptoms	(check w/dr.)	(check w/dr.)	¹∕2 tsp	1 tsp	1 ½ tsp
Dimetapp Elixir	(check w/dr.)	(check w/dr.)	½ tsp	¹∕2 tsp	³⁄4 tsp
Triaminic Syrup (orange)	(CHECK W/GI.)	(check w/ui.)	½ tsp	³ / ₄ tsp	1 tsp
Every 6 hours as needed For congestion and Drainage. May help Some types of coughs			/1 to p	74 65 6	1 65 p
Robitussin or Robitussin PE may Be used for cough every 6 hours as needed	(check w/dr.)	(check w/dr.)	1/4 tsp	½ tsp	3/4 tsp
	Physician's Signature			Date	